

ANALYSIS OF THE $\frac{1}{2}^-$ AND $\frac{3}{2}^-$ HEAVY AND DOUBLY HEAVY BARYON STATES WITH QCD SUM RULES

Zhi-Gang Wang¹

Department of Physics, North China Electric Power University, Baoding 071003, P. R. China

Abstract

In this article, we study the $\frac{1}{2}^-$ and $\frac{3}{2}^-$ heavy and doubly heavy baryon states $\Sigma_Q(\frac{1}{2}^-)$, $\Xi'_Q(\frac{1}{2}^-)$, $\Omega_Q(\frac{1}{2}^-)$, $\Xi_{QQ}(\frac{1}{2}^-)$, $\Omega_{QQ}(\frac{1}{2}^-)$, $\Sigma_Q^*(\frac{3}{2}^-)$, $\Xi_Q^*(\frac{3}{2}^-)$, $\Omega_Q^*(\frac{3}{2}^-)$, $\Xi_{QQ}^*(\frac{3}{2}^-)$ and $\Omega_{QQ}^*(\frac{3}{2}^-)$ by subtracting the contributions from the corresponding $\frac{1}{2}^+$ and $\frac{3}{2}^+$ heavy and doubly heavy baryon states with the QCD sum rules in a systematic way, and make reasonable predictions for their masses.

PACS number: 14.20.Lq, 14.20.Mr

Key words: Heavy baryon states, Doubly heavy baryon states, QCD sum rules

1 Introduction

In the past years, there have been several experimental progresses on the spectroscopy of the heavy baryon states. In 2006, the Babar collaboration reported the first observation of the $\frac{3}{2}^+$ excited singly-charmed baryon state Ω_c^* in the radiative decay $\Omega_c^* \rightarrow \Omega_c \gamma$ at the PEP-II asymmetric energy B -factory at the Stanford linear accelerator center [1]. By now, the $\frac{1}{2}^+$ and $\frac{1}{2}^-$ antitriplet charmed baryon states (Λ_c^+ , Ξ_c^+ , Ξ_c^0) and ($\Lambda_c^+(2595)$, $\Xi_c^+(2790)$, $\Xi_c^0(2790)$), and the $\frac{1}{2}^+$ and $\frac{3}{2}^+$ sextet charmed baryon states (Ω_c , Σ_c , Ξ_c') and (Ω_c^* , Σ_c^* , Ξ_c^*) have been observed [2]. In 2008, the D0 collaboration reported the first observation of the doubly strange baryon state Ω_b^- in the decay channel $\Omega_b^- \rightarrow J/\psi \Omega^-$ with $J/\psi \rightarrow \mu^+ \mu^-$ and $\Omega^- \rightarrow \Lambda K^-$ in the $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV [3]. However, the CDF collaboration did not confirm the measured mass $M_{\Omega_b^-} = (6.165 \pm 0.010 \pm 0.013)$ GeV [4], they determined the mass to be $(6.0544 \pm 0.0068 \pm 0.0009)$ GeV. By now, the S -wave bottom baryon states are far from complete, only the Λ_b , Σ_b , Σ_b^* , Ξ_b , Ω_b have been observed [2]. In 2002, the SELEX collaboration reported the first observation of a signal for the doubly charmed baryon state Ξ_{cc}^+ in the charged decay mode $\Xi_{cc}^+ \rightarrow \Lambda_c^+ K^- \pi^+$ at the charm hadro-production experiment (E781) [5], and confirmed later by the same collaboration in the decay mode $\Xi_{cc}^+ \rightarrow p D^+ K^-$ with measured mass $M_{\Xi} = (3518.9 \pm 0.9)$ MeV [6]. However, the Babar and Belle collaborations have not observed any evidence for the doubly charmed baryon states in e^+e^- annihilations [7, 8], and did not confirm the SELEX observations. Moreover, several new excited charmed baryon states, such as the $\Lambda_c(2765)^+$, $\Lambda_c^+(2880)$, $\Lambda_c^+(2940)$, $\Sigma_c^+(2800)$, $\Xi_c^+(2980)$, $\Xi_c^+(3080)$, $\Xi_c^0(2980)$, $\Xi_c^0(3080)$, have been observed by the Babar, Belle and CLEO collaborations in recent years [9, 10, 11, 12], and re-vivified the interest in the spectroscopy of the charmed baryon states. On the other hand, the large hadron collider (LHC) will provide us with the whole excited bottom baryon states [13]. Therefore it is interesting to calculate the mass spectrum of the excited heavy and doubly heavy baryon states with the QCD sum rules.

There have been several approaches to deal with the heavy and doubly heavy baryon masses, many works focus on the positive-parity baryon states (one can consult Ref.[14] for more literatures); while the works on the negative-parity baryon states are relatively few, for example, the non-relativistic quark model [14, 15], the relativized quark model [16, 17], the relativistic quark model based on a quasipotential approach in QCD [18, 19], the potential approach combined with the QCD sum rules [20], the full QCD sum rules [21, 22, 23], the three-body Faddeev method [24], etc.

The QCD sum rules is a powerful theoretical tool in studying the ground state heavy baryons [25, 26, 27]. In the QCD sum rules, the operator product expansion is used to expand the time-ordered currents into a series of quark and gluon condensates which parameterize the long distance

¹E-mail, wangzgyiti@yahoo.com.cn.

properties of the QCD. Based on the quark-hadron duality, we can obtain copious information about the hadronic parameters at the phenomenological side [25, 26, 27]. There have been several works on the masses of the heavy baryon states with the full QCD sum rules and the QCD sum rules in the heavy quark effective theory (one can consult Ref.[28] for more literatures).

In Ref.[29], Jido et al introduce a novel approach based on the QCD sum rules to separate the contributions of the negative-parity light flavor baryons from the positive-parity light flavor baryons, as the interpolating currents may have non-vanishing couplings to both the negative- and positive-parity baryons [30]. Before the work of Jido et al, Bagan et al take the infinite mass limit for the heavy quarks to separate the contributions of the positive- and negative-parity heavy baryon states unambiguously [31].

In Refs.[28, 32, 33, 34], we follow Ref.[29] and study the masses and pole residues of the $\frac{1}{2}^+$ and $\frac{3}{2}^+$ heavy and doubly heavy baryon states $\Lambda_Q, \Xi_Q, \Sigma_Q, \Xi'_Q, \Omega_Q, \Xi_{QQ}, \Omega_{QQ}, \Sigma_Q^*, \Xi_Q^*, \Omega_Q^*, \Xi_{QQ}^*$ and Ω_{QQ}^* in a systematic way by subtracting the contributions of the corresponding negative parity heavy and doubly heavy baryon states. In this article, we extend our previous works to study the $\frac{1}{2}^-$ and $\frac{3}{2}^-$ heavy and doubly heavy baryon states $\Sigma_Q(\frac{1}{2}^-), \Xi'_Q(\frac{1}{2}^-), \Omega_Q(\frac{1}{2}^-), \Xi_{QQ}(\frac{1}{2}^-), \Omega_{QQ}(\frac{1}{2}^-), \Sigma_Q^*(\frac{3}{2}^-), \Xi_Q^*(\frac{3}{2}^-), \Omega_Q^*(\frac{3}{2}^-), \Xi_{QQ}^*(\frac{3}{2}^-)$ and $\Omega_{QQ}^*(\frac{3}{2}^-)$ by subtracting the contributions from the corresponding $\frac{1}{2}^+$ and $\frac{3}{2}^+$ heavy and doubly heavy baryon states with the full QCD sum rules.

The article is arranged as follows: we derive the QCD sum rules for the masses and the pole residues of the negative parity heavy and doubly heavy baryon states $\Sigma_Q(\frac{1}{2}^-), \Xi'_Q(\frac{1}{2}^-), \Omega_Q(\frac{1}{2}^-), \Xi_{QQ}(\frac{1}{2}^-), \Omega_{QQ}(\frac{1}{2}^-), \Sigma_Q^*(\frac{3}{2}^-), \Xi_Q^*(\frac{3}{2}^-), \Omega_Q^*(\frac{3}{2}^-), \Xi_{QQ}^*(\frac{3}{2}^-)$ and $\Omega_{QQ}^*(\frac{3}{2}^-)$ in Sect.2; in Sect.3, we present the numerical results and discussions; and Sect.4 is reserved for our conclusions.

2 QCD sum rules for the negative parity heavy and doubly heavy baryon states

The ground state quarks have the spin-parity $\frac{1}{2}^+$, two quarks can form a scalar diquark or an axialvector diquark with the spin-parity 0^+ or 1^+ without additional relative P -wave, the diquark then combines with a third quark to form a positive parity baryon. If there exists a relative P -wave (which can be denoted as 1^-) between the diquark and the third quark, we can combine the relative P -wave with the third quark to form an excited quark with the $J^P = \frac{1}{2}^-$ or $\frac{3}{2}^-$ firstly, then combine the negative-parity quark with the positive parity diquark to form a negative parity baryon. We can denote the quarks, diquarks and baryons with their spin-parity (or J^P), and illustrate the process in the following formula explicitly,

$$\begin{aligned} \left[\frac{1}{2}^+ \otimes \frac{1}{2}^+ \right] \otimes \frac{1}{2}^+ &= [0^+ + 1^+] \otimes \frac{1}{2}^+ = \frac{1}{2}^+ + \left[\frac{1}{2}^+ + \frac{3}{2}^+ \right], \\ \left[\frac{1}{2}^+ \otimes \frac{1}{2}^+ \right] \otimes \left[\frac{1}{2}^+ \otimes 1^- \right] &= [0^+ + 1^+] \otimes \left[\frac{1}{2}^- + \frac{3}{2}^- \right], \\ 1^+ \otimes \frac{1}{2}^+ &= \frac{1}{2}^+ + \frac{3}{2}^+, \\ 1^+ \otimes \frac{1}{2}^- &= \frac{1}{2}^- + \frac{3}{2}^-, \end{aligned} \tag{1}$$

the baryons with the 0^+ and 1^+ diquarks are usually called the Λ -type and Σ -type baryons respectively [35]. In the heavy quark limit, the heavy and doubly heavy baryon states can be described by the diquark-quark model [20]. In the quantum field theory, we usually construct the 0^+ and 1^+ diquark currents with the anti-symmetric Dirac matrix $C\gamma_5$ and the symmetric Dirac matrix

$C\gamma_\mu$, respectively. In this article, we construct the Σ -type currents to interpolate the $\frac{1}{2}^\pm$ and $\frac{3}{2}^\pm$ heavy and doubly heavy baryon states.

The $\frac{1}{2}^+$ and $\frac{3}{2}^+$ heavy and doubly heavy baryon states $\Sigma_Q, \Xi'_Q, \Omega_Q, \Xi_{QQ}, \Omega_{QQ}, \Sigma_Q^*, \Xi_Q^*, \Omega_Q^*, \Xi_{QQ}^*$ and Ω_{QQ}^* can be interpolated by the following currents $J^{\Sigma_Q}(x), J^{\Xi'_Q}(x), J^{\Omega_Q}(x), J^{\Xi_{QQ}}(x), J^{\Omega_{QQ}}(x), J^{\Sigma_Q^*}(x), J^{\Xi_Q^*}(x), J^{\Omega_Q^*}(x), J^{\Xi_{QQ}^*}(x)$ and $J^{\Omega_{QQ}^*}(x)$, respectively,

$$\begin{aligned}
J^{\Sigma_Q}(x) &= \epsilon^{ijk} u_i^T(x) C \gamma_\mu d_j(x) \gamma^\mu \gamma_5 Q_k(x), \\
J^{\Xi'_Q}(x) &= \epsilon^{ijk} q_i^T(x) C \gamma_\mu s_j(x) \gamma^\mu \gamma_5 Q_k(x), \\
J^{\Omega_Q}(x) &= \epsilon^{ijk} s_i^T(x) C \gamma_\mu s_j(x) \gamma^\mu \gamma_5 Q_k(x), \\
J^{\Xi_{QQ}}(x) &= \epsilon^{ijk} Q_i^T(x) C \gamma_\mu Q_j(x) \gamma^\mu \gamma_5 q_k(x), \\
J^{\Omega_{QQ}}(x) &= \epsilon^{ijk} Q_i^T(x) C \gamma_\mu Q_j(x) \gamma^\mu \gamma_5 s_k(x), \\
J^{\Sigma_Q^*}(x) &= \epsilon^{ijk} u_i^T(x) C \gamma_\mu d_j(x) Q_k(x), \\
J^{\Xi_Q^*}(x) &= \epsilon^{ijk} q_i^T(x) C \gamma_\mu s_j(x) Q_k(x), \\
J^{\Omega_Q^*}(x) &= \epsilon^{ijk} s_i^T(x) C \gamma_\mu s_j(x) Q_k(x), \\
J^{\Xi_{QQ}^*}(x) &= \epsilon^{ijk} Q_i^T(x) C \gamma_\mu Q_j(x) q_k(x), \\
J^{\Omega_{QQ}^*}(x) &= \epsilon^{ijk} Q_i^T(x) C \gamma_\mu Q_j(x) s_k(x),
\end{aligned} \tag{2}$$

where the Q represents the heavy quarks c and b , the i, j and k are color indexes, and the C is the charge conjugation matrix. In this article, we take the Ioffe-type interpolating currents for the spin $\frac{1}{2}$ baryon states for simplicity [36], the general expression of the spin $\frac{1}{2}$ baryon currents is

$$\eta(x) = \epsilon^{ijk} [\psi_i(x) C \gamma_5 \psi_j(x) \psi_k(x) + t \psi_i(x) C \psi_j(x) \gamma_5 \psi_k(x)],$$

where the $\psi_i(x)$ denote the quarks q, s and Q [27, 31, 37]. If one choose the optimal parameter t , the results can be improved.

The corresponding $\frac{1}{2}^-$ and $\frac{3}{2}^-$ heavy and doubly heavy baryon states can be interpolated by the currents $J^- = i\gamma_5 J^+$ and $J_\mu^- = i\gamma_5 J_\mu^+$ respectively, because multiplying $i\gamma_5$ to the J^+ and J_μ^+ changes the parity of the J^+ and J_μ^+ [29], where the J^+ and J_μ^+ denote the currents interpolating the $\Sigma_Q, \Xi'_Q, \Omega_Q, \Xi_{QQ}, \Omega_{QQ}$ and $\Sigma_Q^*, \Xi_Q^*, \Omega_Q^*, \Xi_{QQ}^*, \Omega_{QQ}^*$, respectively.

The correlation functions $\Pi^\pm(p)$ and $\Pi_{\mu\nu}^\pm(p)$ are defined by

$$\begin{aligned}
\Pi^\pm(p) &= i \int d^4x e^{ip \cdot x} \langle 0 | T \{ J^\pm(x) \bar{J}^\pm(0) \} | 0 \rangle, \\
\Pi_{\mu\nu}^\pm(p) &= i \int d^4x e^{ip \cdot x} \langle 0 | T \{ J_\mu^\pm(x) \bar{J}_\nu^\pm(0) \} | 0 \rangle.
\end{aligned} \tag{3}$$

The currents $J_\pm$ couple to $\frac{1}{2}^\pm$ baryon states $B_\pm$, while the currents $J_\mu^\pm(x)$ couple to both the $\frac{3}{2}^\pm$ baryon states $B_\pm^*$ and the $\frac{1}{2}^\pm$ baryon states $B_\pm$ [30], i.e.

$$\begin{aligned}
\langle 0 | J_+(0) | B_\pm(p) \rangle \langle B_\pm(p) | \bar{J}_+(0) | 0 \rangle &= -\gamma_5 \langle 0 | J_-(0) | B_\pm(p) \rangle \langle B_\pm(p) | \bar{J}_-(0) | 0 \rangle \gamma_5, \\
\langle 0 | J_\mu^+(0) | B_\pm^*(p) \rangle \langle B_\pm^*(p) | \bar{J}_\mu^+(0) | 0 \rangle &= -\gamma_5 \langle 0 | J_\mu^-(0) | B_\pm^*(p) \rangle \langle B_\pm^*(p) | \bar{J}_\mu^-(0) | 0 \rangle \gamma_5, \\
\langle 0 | J_\mu^+(0) | B_\pm(p) \rangle \langle B_\pm(p) | \bar{J}_\mu^+(0) | 0 \rangle &= -\gamma_5 \langle 0 | J_\mu^-(0) | B_\pm(p) \rangle \langle B_\pm(p) | \bar{J}_\mu^-(0) | 0 \rangle \gamma_5,
\end{aligned} \tag{4}$$

where

$$\begin{aligned}
\langle 0 | J^\pm(0) | B_\pm(p) \rangle &= \lambda_\pm U(p, s), \\
\langle 0 | J_\mu^\pm(0) | B_\pm^*(p) \rangle &= \lambda_\pm U_\mu(p, s), \\
\langle 0 | J_\mu^\pm(0) | B_\mp(p) \rangle &= \lambda_\mp \left(\gamma_\mu - 4 \frac{p_\mu}{M_\mp} \right) U(p, s),
\end{aligned} \tag{5}$$

the $\lambda_{\pm}$ are the pole residues and the $M_{\pm}$ are the masses, and the spinor $U(p, s)$ satisfies the usual Dirac equation $(\not{p} - M_{\pm})U(p) = 0$.

The $\Pi^{\pm}(p)$ and $\Pi_{\mu\nu}^{\pm}(p)$ have the following relation

$$\begin{aligned}\Pi^{-}(p) &= -\gamma_5 \Pi^{+}(p) \gamma_5, \\ \Pi_{\mu\nu}^{-}(p) &= -\gamma_5 \Pi_{\mu\nu}^{+}(p) \gamma_5.\end{aligned}\tag{6}$$

We insert a complete set of intermediate baryon states with the same quantum numbers as the current operators $J^{\pm}(x)$ and $J_{\mu}^{\pm}(x)$ into the correlation functions $\Pi^{-}(p)$ and $\Pi_{\mu\nu}^{-}(p)$ to obtain the hadronic representation [25, 26]. After isolating the pole terms of the lowest states of the heavy and doubly heavy baryons, we obtain the following results [29]:

$$\begin{aligned}\Pi^{-}(p) &= \lambda_-^2 \frac{\not{p} - M_-}{M_-^2 - p^2} + \lambda_+^2 \frac{\not{p} + M_+}{M_+^2 - p^2} + \cdots, \\ \Pi_{\mu\nu}^{-}(p) &= -\lambda_-^2 \frac{\not{p} - M_-}{M_-^2 - p^2} g_{\mu\nu} - \lambda_+^2 \frac{\not{p} + M_+}{M_+^2 - p^2} g_{\mu\nu} + \cdots, \\ &= -\Pi_{-}(p) g_{\mu\nu} + \cdots,\end{aligned}\tag{7}$$

where the $M_{\pm}$ are the masses of the lowest states with the parity $\pm$ respectively, and the $\lambda_{\pm}$ are the corresponding pole residues (or couplings). In this article, we choose the tensor structure $g_{\mu\nu}$ for analysis, the $\frac{1}{2}^{\pm}$ baryon states have no contaminations.

If we take $\vec{p} = 0$ (here we use the $\Pi(p)$ to denote the $\Pi^{-}(p)$ and $\Pi_{-}(p)$ together), then

$$\begin{aligned}\lim_{\epsilon \rightarrow 0} \frac{\text{Im}\Pi(p_0 + i\epsilon)}{\pi} &= \lambda_-^2 \frac{\gamma_0 - 1}{2} \delta(p_0 - M_-) + \lambda_+^2 \frac{\gamma_0 + 1}{2} \delta(p_0 - M_+) + \cdots \\ &= \gamma_0 A(p_0) - B(p_0) + \cdots,\end{aligned}\tag{8}$$

where

$$\begin{aligned}A(p_0) &= \frac{1}{2} [\lambda_-^2 \delta(p_0 - M_-) + \lambda_+^2 \delta(p_0 - M_+)], \\ B(p_0) &= \frac{1}{2} [\lambda_-^2 \delta(p_0 - M_-) - \lambda_+^2 \delta(p_0 - M_+)],\end{aligned}\tag{9}$$

the $A(p_0) + B(p_0)$ and $A(p_0) - B(p_0)$ contain the contributions from the negative-parity baryon states and positive-parity baryon states, respectively.

We calculate the light quark parts of the correlation functions $\Pi^{-}(p)$ and $\Pi_{\mu\nu}^{-}(p)$ in the coordinate space and use the momentum space expression for the heavy quark propagators, i.e. we take

$$\begin{aligned}S_{ij}(x) &= \frac{i\delta_{ij}\not{x}}{2\pi^2 x^4} - \frac{\delta_{ij}m_s}{4\pi^2 x^2} - \frac{\delta_{ij}}{12}\langle\bar{s}s\rangle + \frac{i\delta_{ij}}{48}m_s\langle\bar{s}s\rangle\not{x} \\ &\quad - \frac{i}{32\pi^2 x^2}G_{\mu\nu}^{ij}(x)[\not{x}\sigma^{\mu\nu} + \sigma^{\mu\nu}\not{x}] + \cdots, \\ S_Q^{ij}(x) &= \frac{i}{(2\pi)^4} \int d^4k e^{-ik\cdot x} \left\{ \frac{\delta_{ij}}{\not{k} - m_Q} - \frac{g_s G_{ij}^{\alpha\beta}}{4} \frac{\sigma_{\alpha\beta}(\not{k} + m_Q) + (\not{k} + m_Q)\sigma_{\alpha\beta}}{(k^2 - m_Q^2)^2} \right. \\ &\quad \left. + \frac{\pi^2}{3} \langle \frac{\alpha_s GG}{\pi} \rangle \delta_{ij} m_Q \frac{k^2 + m_Q \not{k}}{(k^2 - m_Q^2)^4} + \cdots \right\},\end{aligned}\tag{10}$$

where $\langle \frac{\alpha_s GG}{\pi} \rangle = \langle \frac{\alpha_s G_{\alpha\beta} G^{\alpha\beta}}{\pi} \rangle$, then resort to the Fourier integral to transform the light quark parts into the momentum space in D dimensions, take $\vec{p} = 0$, and use the dispersion relation to obtain the spectral densities $\rho^A(p_0)$ and $\rho^B(p_0)$ (which correspond to the tensor structures γ_0 and

1 respectively) at the level of quark-gluon degrees of freedom. Finally we introduce the weight functions $\exp\left[-\frac{p_0^2}{T^2}\right]$, $p_0^2 \exp\left[-\frac{p_0^2}{T^2}\right]$, and obtain the following sum rules,

$$\lambda_-^2 \exp\left[-\frac{M_-^2}{T^2}\right] = \int_{\Delta}^{\sqrt{s_0}} dp_0 [\rho^A(p_0) + \rho^B(p_0)] \exp\left[-\frac{p_0^2}{T^2}\right], \quad (11)$$

$$\lambda_-^2 M_-^2 \exp\left[-\frac{M_-^2}{T^2}\right] = \int_{\Delta}^{\sqrt{s_0}} dp_0 [\rho^A(p_0) + \rho^B(p_0)] p_0^2 \exp\left[-\frac{p_0^2}{T^2}\right], \quad (12)$$

where the s_0 are the threshold parameters, the T^2 are the Borel parameters, and $\Delta = 2m_Q + m_s$, $2m_Q$, $m_Q + 2m_s$, $m_Q + m_s$ and m_Q in the channels $\Omega_{QQ}^{(*)}$, $\Xi_{QQ}^{(*)}$, $\Omega_Q^{(*)}$, $\Xi_Q^{(*)}$ and $\Sigma_Q^{(*)}$, respectively. The spectral densities $\rho^A(p_0)$ and $\rho^B(p_0)$ at the level of quark-gluon degrees of freedom are given explicitly in the Appendix.

3 Numerical results and discussions

The input parameters are taken to be the standard values $\langle \bar{q}q \rangle = -(0.24 \pm 0.01 \text{ GeV})^3$, $\langle \bar{s}s \rangle = (0.8 \pm 0.2) \langle \bar{q}q \rangle$, $\langle \bar{q}g_s \sigma G q \rangle = m_0^2 \langle \bar{q}q \rangle$, $\langle \bar{s}g_s \sigma G s \rangle = m_0^2 \langle \bar{s}s \rangle$, $m_0^2 = (0.8 \pm 0.2) \text{ GeV}^2$ [38, 39], $\langle \frac{\alpha_s GG}{\pi} \rangle = (0.012 \pm 0.004) \text{ GeV}^4$ [39], $m_s = (0.14 \pm 0.01) \text{ GeV}$, $m_c = (1.35 \pm 0.10) \text{ GeV}$ and $m_b = (4.7 \pm 0.1) \text{ GeV}$ [2] at the energy scale $\mu^2 = 1 \text{ GeV}^2$.

The value of the gluon condensate $\langle \frac{\alpha_s GG}{\pi} \rangle$ has been updated from time to time, and changes greatly [27]. At the present case, the gluon condensate makes tiny contribution, the updated value $\langle \frac{\alpha_s GG}{\pi} \rangle = (0.023 \pm 0.003) \text{ GeV}^4$ [27] and the standard value $\langle \frac{\alpha_s GG}{\pi} \rangle = (0.012 \pm 0.004) \text{ GeV}^4$ [39] lead to a tiny difference and can be neglected safely.

For the light quark masses, we take the approximation $m_u = m_d \approx 0$. The values listed in the Review of Particle Physics are $m_u = 2.49_{-0.79}^{+0.81} \text{ MeV}$ and $m_d = 5.05_{-0.95}^{+0.75} \text{ MeV}$ at the energy scale $\mu = 2 \text{ GeV}$ [2]. The values $m_q = \frac{m_u + m_d}{2} = 5.6 \text{ MeV}$ [27] and $m_q = 0$ lead to a difference less than 2 MeV for masses of the Qqs , Qqq' and QQq baryon states, we can neglect the u and d quark masses safely.

The Q -quark masses appearing in the perturbative terms are usually taken to be the pole masses in the QCD sum rules, while the choice of the m_Q in the leading-order coefficients of the higher-dimensional terms is arbitrary [27, 40]. The $\overline{MS}$ mass $m_c(m_c^2)$ relates with the pole mass $\hat{m}_c$ through the relation $m_c(m_c^2) = \hat{m}_c \left[1 + \frac{C_F \alpha_s(m_c^2)}{\pi} + \dots\right]^{-1}$. In this article, we take the approximation $m_c(m_c^2) \approx \hat{m}_c$ without the α_s corrections for consistency. The value listed in the Review of Particle Physics is $m_c(m_c^2) = 1.27_{-0.09}^{+0.07} \text{ GeV}$ [2], it is reasonable to take $\hat{m}_c = m_c(1 \text{ GeV}^2) = (1.35 \pm 0.10) \text{ GeV}$. For the b quark, the $\overline{MS}$ mass is $m_b(m_b^2) = 4.19_{-0.06}^{+0.18} \text{ GeV}$ [2], the gap between the energy scale $\mu = 4.2 \text{ GeV}$ and 1 GeV is rather large, the approximation $\hat{m}_b \approx m_b(m_b^2) \approx m_b(1 \text{ GeV}^2)$ seems rather crude. It would be better to understand the heavy quark masses m_c and m_b we take at the energy scale $\mu^2 = 1 \text{ GeV}^2$ as the effective quark masses (or just the mass parameters). Our previous works on the mass spectrum of the $\frac{1}{2}^+$ and $\frac{3}{2}^+$ heavy and doubly heavy baryon states and the $\frac{1}{2}^-$ antitriplet heavy baryon states indicate such parameters can lead to satisfactory results [28, 32, 33, 34].

In Ref.[21], Bagan et al observe that the contribution from the mixed condensates has a term that behaves like $\frac{1}{v^3}$ (the v is the heavy quark velocity) for the doubly heavy baryon states, which indicates a coulombic correction and requires a complete treatment of the nonrelativistic coulombic corrections. In this article, we take (almost) the same pole contributions and analogous convergent behaviors in the operator product expansion as our previous works [33, 34], where the positive-parity doubly heavy baryon states were studied with the vacuum condensates up to dimension 4, and neglect the contributions from the mixed condensates for consistency. We also present the

results with the mixed condensates for comparison, see Table 1. For the $\frac{1}{2}^-$ doubly heavy baryon states, there are cancellations among the mixed condensates which originate from the diagrams where the gluons emitted from the two heavy quarks and the light quark respectively and then absorbed by the light quark. The net contributions are minor and can be neglected safely. On the other hand, for the $\frac{3}{2}^-$ doubly heavy baryon states, there are only contributions from the mixed condensates which originate from the diagrams where the gluons emitted from the light quark and then absorbed by the light quark itself. There are no cancellations, the inclusion of the mixed condensates can result in masses about $(30 - 50)$ MeV larger than the corresponding ones without them, and the uncertainties of the masses originate from the parameter $m_0^2 = (0.8 \pm 0.2) \text{ GeV}^2$ are less than 10 MeV.

In calculations, we neglect the contributions from the perturbative corrections $\mathcal{O}(\alpha_s^n)$. Those perturbative corrections can be taken into account in the leading logarithmic approximations through anomalous dimension factors. After the Borel transform, the effects of those corrections are to multiply each term on the operator product expansion side by the factor, $\left[\frac{\alpha_s(T^2)}{\alpha_s(\mu^2)}\right]^{2\Gamma_J - \Gamma_{\mathcal{O}_n}}$, where the Γ_J is the anomalous dimension of the interpolating current $J(x)$ and the $\Gamma_{\mathcal{O}_n}$ is the anomalous dimension of the local operator $\mathcal{O}_n(0)$. We carry out the operator product expansion at a special energy scale $\mu^2 = 1 \text{ GeV}^2$, and set the factor $\left[\frac{\alpha_s(T^2)}{\alpha_s(\mu^2)}\right]^{2\Gamma_J - \Gamma_{\mathcal{O}_n}} \approx 1$, such an approximation maybe result in some scale dependence and weaken the prediction ability. In this article, we study the $\frac{1}{2}^-$ and $\frac{3}{2}^-$ heavy and doubly heavy baryon states systemically, the predictions are still robust as we take the analogous criteria in those sum rules.

In Refs.[28, 32, 33, 34], we study the masses of the $\frac{1}{2}^\pm$ antitriplet heavy baryon states $\Lambda_Q(\frac{1}{2}^\pm)$, $\Xi_Q(\frac{1}{2}^\pm)$, and the $\frac{1}{2}^+$ and $\frac{3}{2}^+$ heavy and doubly heavy baryon states Σ_Q , Ξ'_Q , Ω_Q , Ξ_{QQ} , Ω_{QQ} , Σ_Q^* , Ξ_Q^* , Ω_Q^* , Ξ_{QQ}^* and Ω_{QQ}^* systematically by subtracting the contributions of the corresponding negative (or positive) parity heavy and doubly heavy baryon states, and obtain satisfactory results. In calculations, we take the values $\sqrt{s_0} - M_{\text{gr}} \approx (0.7 - 0.8) \text{ GeV}$ and the pole contributions $\approx (45 - 80)\%$ and $(45 - 70)\%$ in the charmed channels and the bottom channels respectively, where the s_0 stands for the central values of the threshold parameters, and the M_{gr} stands for the experimental values of the heavy and doubly heavy baryon states $\Lambda_c(\frac{1}{2}^\pm)$, $\Xi_c(\frac{1}{2}^\pm)$, Λ_b , Ξ_b , Σ_c , Ξ'_c , Ω_c , Σ_b , Ω_b , Σ_c^* , Ξ_c^* , Ω_c^* , Σ_b^* , Ω_{cc} listed in the Review of Particle Physics [2], here we take the average value of the mass of the Ω_b from the experimental data of the D0 and CDF collaborations [3, 4].

In the conventional QCD sum rules [25, 26], there are two criteria (pole dominance and convergence of the operator product expansion) for choosing the Borel parameters T^2 and threshold parameters s_0 . We can borrow some ideas from our previous works [28, 32, 33, 34], and take the values $\sqrt{s_0} = M_{\text{gr}} + 0.7 \text{ GeV}$ as guides (where the M_{gr} denotes the ground state masses of the negative parity heavy and doubly heavy baryon states from the non-relativistic quark model [14]), and impose the two criteria on the heavy and doubly heavy baryon states to choose the Borel parameters T^2 and threshold parameters s_0 . If we take the contributions of the pole terms to be $(45 - 80)\%$ and $(45 - 70)\%$ in the charmed channels and the bottom channels respectively (for the central values of the threshold parameters, the pole contributions are larger than or equal 50%), and the contributions of the perturbative terms to be $(50 - 90)\%$ in the operator product expansion as in our previous works [28, 32, 33, 34], we can obtain approximately uniform Borel windows. In this article, we take the uniform Borel windows $T_{\text{max}}^2 - T_{\text{min}}^2 = 1.0 \text{ GeV}^2$, 1.5 GeV^2 and 2.0 GeV^2 in the singly heavy baryon channels, the doubly charmed baryon channels and the doubly bottom baryon channels, respectively; the corresponding threshold parameters are presented in Table 1. The two criteria of the QCD sum rules are fully satisfied [25, 26]. If we take the values $T^2 < T_{\text{min}}^2$, the resulting masses and pole residues change quickly with variations of the Borel parameters, and no satisfactory Borel plateaus can be obtained. On the other hand, if we take the values $T^2 > T_{\text{max}}^2$, the pole contributions are reduced remarkably, we have to postpone the threshold parameters s_0 to larger values, the contributions from the higher resonances and continuum states

may be included in, which will weaken the prediction ability. Although with fine-tuning, we can obtain slightly larger Borel windows, the predictions cannot be improved remarkably, we prefer the uniform Borel windows 1 GeV^2 , 1.5 GeV^2 and 2.0 GeV^2 in the due channels.

In this article, we take uniform uncertainties for the threshold parameters, $\delta\sqrt{s_0} = \pm 0.1 \text{ GeV}$. In calculations, we observe that the predicted masses are not sensitive to the threshold parameters in the Borel windows, although they increase with the threshold parameters, the higher resonances and continuum states are suppressed by the factor $\exp\left[-\frac{p_0^2}{T^2}\right] \Theta(p_0^2 - s_0) \leq e^{-1}$. In fact, we can take larger uncertainties, $\delta\sqrt{s_0} = \pm 0.2 \text{ GeV}$, which can enlarge the uncertainties of the masses about $(0.01 - 0.03) \text{ GeV}$. For example, if we take the larger uncertainties $\delta\sqrt{s_0} = \pm 0.2 \text{ GeV}$ and the same pole contributions, the Borel parameters in the channels Ω_c , Ω_c^* , Ξ_c' and Ξ_c^* should be taken as $T^2 \approx (2.9 - 3.5) \text{ GeV}^2$, $(2.9 - 3.5) \text{ GeV}^2$, $(2.7 - 3.3) \text{ GeV}^2$ and $(2.7 - 3.3) \text{ GeV}^2$, respectively, i.e. the Borel windows shrink, the predicted masses are $M = (2.98 \pm 0.18) \text{ GeV}$, $(2.98 \pm 0.18) \text{ GeV}$, $(2.87 \pm 0.18) \text{ GeV}$ and $(2.87 \pm 0.19) \text{ GeV}$, respectively; the uncertainties are enlarged by $(0.01 - 0.03) \text{ GeV}$.

Taking into account all uncertainties of the relevant parameters, we can obtain the values of the masses and pole residues of the $\frac{1}{2}^-$ and $\frac{3}{2}^-$ heavy and doubly heavy baryon states $\Sigma_Q(\frac{1}{2}^-)$, $\Xi_Q'(\frac{1}{2}^-)$, $\Omega_Q(\frac{1}{2}^-)$, $\Xi_{QQ}(\frac{1}{2}^-)$, $\Omega_{QQ}(\frac{1}{2}^-)$, $\Sigma_Q^*(\frac{3}{2}^-)$, $\Xi_Q^*(\frac{3}{2}^-)$, $\Omega_Q^*(\frac{3}{2}^-)$, $\Xi_{QQ}^*(\frac{3}{2}^-)$ and $\Omega_{QQ}^*(\frac{3}{2}^-)$, which are shown in Figs.1-4 and Table 1. In this article, we calculate the uncertainties δ with the formula

$$\delta = \sqrt{\sum_i \left(\frac{\partial f}{\partial x_i}\right)^2 \big|_{x_i=\bar{x}_i} (x_i - \bar{x}_i)^2}, \quad (13)$$

where the f denotes the hadron mass M_- and the pole residue λ_- , the x_i denotes the input QCD parameters m_c , m_b , $\langle\bar{q}q\rangle$, $\langle\bar{s}s\rangle$, $\dots$, and the threshold parameter s_0 and Borel parameter M^2 . As the partial derivatives $\frac{\partial f}{\partial x_i}$ are difficult to carry out analytically, we take the approximation $\left(\frac{\partial f}{\partial x_i}\right)^2 (x_i - \bar{x}_i)^2 \approx [f(\bar{x}_i \pm \Delta x_i) - f(\bar{x}_i)]^2$ in the numerical calculations.

In Table 1, we also present the results from the non-relativistic quark model [14] and the relativistic quark model based on a quasipotential approach in QCD [18, 19], the present predictions are in good agreement with those values within uncertainties.

The charmed partners of the Λ baryon Λ_c^+ , $\Lambda_c^+(2595)$, $\Lambda_c^+(2625)$, $\Lambda_c^+(2765)$ (or $\Sigma_c^+(2765)$), $\Lambda_c^+(2880)$ and $\Lambda_c^+(2940)$ have been observed, the quantum numbers listed in the Review of Particle Physics are $J^P = \frac{1}{2}^+$, $\frac{1}{2}^-$, $\frac{3}{2}^-$, $?$, $\frac{5}{2}^+$ and $?$, respectively [2]. The flux-tube model favors to identify those charmed baryon states with the spin-parity $\frac{1}{2}^+$, $\frac{1}{2}^-$, $\frac{3}{2}^-$, $\frac{3}{2}^+$, $\frac{5}{2}^+$ and $\frac{5}{2}^-$, respectively [41]. In Ref.[32], we study the mass spectrum of the $\frac{1}{2}^\pm$ flavor antitriplet heavy baryon states (Λ_c^+ , Ξ_c^+ , Ξ_c^0) and (Λ_b^0 , Ξ_b^0 , Ξ_b^-) by subtracting the contributions from the corresponding $\frac{1}{2}^\mp$ heavy baryon states with the QCD sum rules, the predictions $M_{\Lambda_c} = (2.26 \pm 0.27) \text{ GeV}$ and $M_{\Lambda_c(2595)} = (2.61 \pm 0.21) \text{ GeV}$ are in good agreement with the experimental data.

The Ξ_c baryon states Ξ_c , Ξ_c' , $\Xi_c(2645)$, $\Xi_c(2790)$, $\Xi_c(2815)$, $\Xi_c(2980)$, $\Xi_c(3055)$, $\Xi_c(3080)$ and $\Xi_c(3123)$ listed in the Review of Particle Physics have the spin-parity $\frac{1}{2}^+$, $\frac{1}{2}^+$, $\frac{3}{2}^+$, $\frac{1}{2}^-$, $\frac{3}{2}^-$, $?$, $?$, $?$ and $?$, respectively [2]. The $\Xi_c(2790)$ and $\Xi_c(2815)$ with the spin-parity $\frac{1}{2}^-$ and $\frac{3}{2}^-$ respectively can be interpreted as the charmed-strange analogues of the $\Lambda_c^+(2595)$ and $\Lambda_c^+(2625)$, or of the $\Lambda(1405)$ and $\Lambda(1520)$; they are flavor antitriplet or Λ -type heavy baryon states [14]. Our predictions $M_{\Xi_c} = (2.44 \pm 0.23) \text{ GeV}$, $M_{\Xi_c'} = (2.56 \pm 0.22) \text{ GeV}$, $M_{\Xi_c(2645)} = (2.65 \pm 0.20) \text{ GeV}$ and $M_{\Xi_c(2790)} = (2.76 \pm 0.18) \text{ GeV}$ are in good agreement with the experimental data [28, 32, 34]. In the non-relativistic quark model [14] and the relativistic quark model based on a quasipotential approach in QCD [18, 19], the $\Xi_c(2815)$ is taken as the Λ -type baryon state. Although the coupling of an Σ -type interpolating current to a Λ -type baryon state is very small, the present prediction $M_{\Xi_c^*} = (2.86 \pm 0.17) \text{ GeV}$ from the Σ -type interpolating current is compatible with the mass of the $\Xi_c(2815)$, an Σ -type $\Xi_c(2815)$ with the spin-parity $\frac{3}{2}^-$ is not excluded. The $\Xi_c(2980)$, $\Xi_c(3055)$,

	$T^2(\text{GeV}^2)$	$\sqrt{s_0}(\text{GeV})$	$M(\text{GeV})$	$\lambda(\text{GeV}^3)$	Ref.[14]	Refs.[18, 19]
$\Sigma_c(\frac{1}{2}^-)$	2.3 – 3.3	3.5 ± 0.1	2.74 ± 0.20	0.071 ± 0.019	2.748	2.795
$\Xi'_c(\frac{1}{2}^-)$	2.5 – 3.5	3.6 ± 0.1	2.87 ± 0.17	0.084 ± 0.019	2.859	2.928
$\Omega_c(\frac{1}{2}^-)$	2.7 – 3.7	3.7 ± 0.1	2.98 ± 0.16	0.136 ± 0.027	2.977	3.020
$\Sigma_b(\frac{1}{2}^-)$	4.9 – 5.9	6.8 ± 0.1	6.00 ± 0.18	0.085 ± 0.022	6.099	6.108
$\Xi'_b(\frac{1}{2}^-)$	5.2 – 6.2	6.9 ± 0.1	6.14 ± 0.15	0.103 ± 0.024	6.192	6.238
$\Omega_b(\frac{1}{2}^-)$	5.5 – 6.5	7.0 ± 0.1	6.27 ± 0.14	0.173 ± 0.035	6.301	6.352
$\Xi_{cc}(\frac{1}{2}^-)$	3.1 – 4.6	4.5 ± 0.1	3.77 ± 0.18	0.159 ± 0.037	3.910	3.838
$\widehat{\Xi}_{cc}(\frac{1}{2}^-)$	3.1 – 4.6	4.5 ± 0.1	3.78 ± 0.18	0.160 ± 0.037	3.910	3.838
$\Omega_{cc}(\frac{1}{2}^-)$	3.4 – 4.9	4.6 ± 0.1	3.91 ± 0.14	0.192 ± 0.041	4.046	4.002
$\widehat{\Omega}_{cc}(\frac{1}{2}^-)$	3.4 – 4.9	4.6 ± 0.1	3.92 ± 0.16	0.192 ± 0.041	4.046	4.002
$\Xi_{bb}(\frac{1}{2}^-)$	8.8 – 10.8	11.1 ± 0.1	10.38 ± 0.15	0.364 ± 0.088	10.493	10.632
$\widehat{\Xi}_{bb}(\frac{1}{2}^-)$	8.8 – 10.8	11.1 ± 0.1	10.39 ± 0.15	0.365 ± 0.089	10.493	10.632
$\Omega_{bb}(\frac{1}{2}^-)$	9.1 – 11.1	11.2 ± 0.1	10.53 ± 0.15	0.443 ± 0.101	10.616	10.771
$\widehat{\Omega}_{bb}(\frac{1}{2}^-)$	9.1 – 11.1	11.2 ± 0.1	10.53 ± 0.15	0.444 ± 0.101	10.616	10.771
$\Sigma_c^*(\frac{3}{2}^-)$	2.4 – 3.4	3.5 ± 0.1	2.74 ± 0.20	0.037 ± 0.009	2.763	2.761
$\Xi_c^*(\frac{3}{2}^-)$	2.6 – 3.6	3.6 ± 0.1	2.86 ± 0.17	0.045 ± 0.009	2.871	2.900
$\Omega_c^*(\frac{3}{2}^-)$	2.8 – 3.8	3.7 ± 0.1	2.98 ± 0.16	0.072 ± 0.013	2.986	2.998
$\Sigma_b^*(\frac{3}{2}^-)$	5.0 – 6.0	6.8 ± 0.1	6.00 ± 0.18	0.047 ± 0.012	6.101	6.076
$\Xi_b^*(\frac{3}{2}^-)$	5.3 – 6.3	6.9 ± 0.1	6.14 ± 0.16	0.054 ± 0.013	6.194	6.212
$\Omega_b^*(\frac{3}{2}^-)$	5.6 – 6.6	7.0 ± 0.1	6.26 ± 0.15	0.095 ± 0.019	6.304	6.330
$\Xi_{cc}^*(\frac{3}{2}^-)$	3.3 – 4.8	4.5 ± 0.1	3.77 ± 0.17	0.087 ± 0.019	3.921	3.959
$\widehat{\Xi}_{cc}^*(\frac{3}{2}^-)$	3.5 – 5.0	4.6 ± 0.1	3.80 ± 0.18	0.095 ± 0.020	3.921	3.959
$\Omega_{cc}^*(\frac{3}{2}^-)$	3.6 – 5.1	4.6 ± 0.1	3.91 ± 0.16	0.105 ± 0.020	4.052	4.102
$\widehat{\Omega}_{cc}^*(\frac{3}{2}^-)$	3.9 – 5.4	4.7 ± 0.1	3.96 ± 0.16	0.116 ± 0.022	4.052	4.102
$\Xi_{bb}^*(\frac{3}{2}^-)$	9.0 – 11.0	11.1 ± 0.1	10.39 ± 0.15	0.206 ± 0.049	10.495	10.647
$\widehat{\Xi}_{bb}^*(\frac{3}{2}^-)$	9.3 – 11.3	11.2 ± 0.1	10.43 ± 0.15	0.227 ± 0.052	10.495	10.647
$\Omega_{bb}^*(\frac{3}{2}^-)$	9.3 – 11.3	11.2 ± 0.1	10.52 ± 0.15	0.251 ± 0.056	10.619	10.785
$\widehat{\Omega}_{bb}^*(\frac{3}{2}^-)$	9.6 – 11.6	11.3 ± 0.1	10.57 ± 0.15	0.275 ± 0.059	10.619	10.785

Table 1: The masses and pole residues of the $\frac{1}{2}^-$ and $\frac{3}{2}^-$ heavy and doubly heavy baryon states, we also present the predictions for the masses from some quark models for comparison. The wide-hat $\widehat{}$ denotes that the mixed condensates are taken into account.




Figure 1: The masses of the $\frac{1}{2}^-$ heavy and doubly heavy baryon states, the A , B , C , D , E , F , G , H , I and J correspond to the channels $\Sigma_c(\frac{1}{2}^-)$, $\Xi'_c(\frac{1}{2}^-)$, $\Omega_c(\frac{1}{2}^-)$, $\Sigma_b(\frac{1}{2}^-)$, $\Xi'_b(\frac{1}{2}^-)$, $\Omega_b(\frac{1}{2}^-)$, $\Xi_{cc}(\frac{1}{2}^-)$, $\Omega_{cc}(\frac{1}{2}^-)$, $\Xi_{bb}(\frac{1}{2}^-)$ and $\Omega_{bb}(\frac{1}{2}^-)$, respectively.




Figure 2: The masses of the $\frac{3}{2}^-$ heavy and doubly heavy baryon states, the A , B , C , D , E , F , G , H , I and J correspond to the channels $\Sigma_c^*(\frac{3}{2}^-)$, $\Xi_c^*(\frac{3}{2}^-)$, $\Omega_c^*(\frac{3}{2}^-)$, $\Sigma_b^*(\frac{3}{2}^-)$, $\Xi_b^*(\frac{3}{2}^-)$, $\Omega_b^*(\frac{3}{2}^-)$, $\Xi_{cc}^*(\frac{3}{2}^-)$, $\Omega_{cc}^*(\frac{3}{2}^-)$, $\Xi_{bb}^*(\frac{3}{2}^-)$ and $\Omega_{bb}^*(\frac{3}{2}^-)$, respectively.




Figure 3: The pole residues of the $\frac{1}{2}^-$ heavy and doubly heavy baryon states, the A , B , C , D , E , F , G , H , I and J correspond to the channels $\Sigma_c(\frac{1}{2}^-)$, $\Xi'_c(\frac{1}{2}^-)$, $\Omega_c(\frac{1}{2}^-)$, $\Sigma_b(\frac{1}{2}^-)$, $\Xi'_b(\frac{1}{2}^-)$, $\Omega_b(\frac{1}{2}^-)$, $\Xi_{cc}(\frac{1}{2}^-)$, $\Omega_{cc}(\frac{1}{2}^-)$, $\Xi_{bb}(\frac{1}{2}^-)$ and $\Omega_{bb}(\frac{1}{2}^-)$, respectively.




Figure 4: The pole residues of the $\frac{3}{2}^-$ heavy and doubly heavy baryon states, the A , B , C , D , E , F , G , H , I and J correspond to the channels $\Sigma_c^*(\frac{3}{2}^-)$, $\Xi_c^*(\frac{3}{2}^-)$, $\Omega_c^*(\frac{3}{2}^-)$, $\Sigma_b^*(\frac{3}{2}^-)$, $\Xi_b^*(\frac{3}{2}^-)$, $\Omega_b^*(\frac{3}{2}^-)$, $\Xi_{cc}^*(\frac{3}{2}^-)$, $\Omega_{cc}^*(\frac{3}{2}^-)$, $\Xi_{bb}^*(\frac{3}{2}^-)$ and $\Omega_{bb}^*(\frac{3}{2}^-)$, respectively.

$\Xi_c(3080)$ and $\Xi_c(3123)$ are unlikely the $\frac{3}{2}^-$ state due to their large masses. The $\Xi_c(3080)$ is tentatively identified as the strange partner of the $\Lambda_c(2880)$ with the spin-parity $\frac{5}{2}^+$ [10, 42], the $\Xi_c(2980)$ maybe have spin-parity $\frac{1}{2}^+$ [18, 42] or $\frac{3}{2}^+$ [10], the $\Xi_c(3055)$ and $\Xi_c(3123)$ can be interpreted as the second orbital ($1D$) excitations of the Ξ_c with $\frac{5}{2}^+$ containing scalar and axial-vector diquarks, respectively [18].

Only three Σ_c baryon states $\Sigma_c(2455)$, $\Sigma_c(2520)$ and $\Sigma_c(2800)$ have been observed, the quantum numbers listed in the Review of Particle Physics are $\frac{1}{2}^+$, $\frac{3}{2}^+$ and ?, respectively [2]. The possible spin-parity of the $\Sigma_c(2800)$ is $\frac{1}{2}^-$, $\frac{3}{2}^-$ and $\frac{5}{2}^-$ [14], or $\frac{3}{2}^-$ [10, 24, 42]. In Refs.[28, 33, 34], we study the $\frac{1}{2}^+$ and $\frac{3}{2}^+$ heavy and doubly heavy baryon states by subtracting the contributions from the corresponding $\frac{1}{2}^-$ and $\frac{3}{2}^-$ heavy and doubly heavy baryon states with the QCD sum rules, the predictions $M_{\Sigma_c(2455)} = (2.40 \pm 0.26) \text{ GeV}$ and $M_{\Sigma_c(2520)} = (2.48 \pm 0.25) \text{ GeV}$ are in good agreement with the experimental data. If we tentatively identify the $\Sigma_c(2800)$ as the $\frac{3}{2}^-$ state, the present prediction $M_{\Sigma_c^*} = (2.74 \pm 0.20) \text{ GeV}$ with the same methods as Refs.[28, 33, 34] is very good.

The properties of the charmed and doubly charmed baryon states would be studied at the BESIII and PANDA [43, 44], where the charmed baryon states are copiously produced at the e^+e^- and $p\bar{p}$ collisions. The LHCb is a dedicated b and c -physics precision experiment at the LHC. The LHC will be the world's most copious source of the b hadrons, and a complete spectrum of the b hadrons will be available through gluon fusion. In proton-proton collisions at $\sqrt{s} = 14 \text{ TeV}$, the $b\bar{b}$ cross section is expected to be $\sim 500 \mu\text{b}$ producing 10^{12} $b\bar{b}$ pairs in a standard year of running at the LHCb operational luminosity of $2 \times 10^{32} \text{ cm}^{-2} \text{ sec}^{-1}$ [13]. The present predictions for the masses of the $\frac{1}{2}^-$ and $\frac{3}{2}^-$ heavy and doubly heavy baryon states can be confronted with the experimental data in the future at the BESIII, PANDA and LHCb.

4 Conclusion

In this article, we study $\frac{1}{2}^-$ and $\frac{3}{2}^-$ heavy and doubly heavy baryon states $\Sigma_Q(\frac{1}{2}^-)$, $\Xi'_Q(\frac{1}{2}^-)$, $\Omega_Q(\frac{1}{2}^-)$, $\Xi_{QQ}(\frac{1}{2}^-)$, $\Omega_{QQ}(\frac{1}{2}^-)$, $\Sigma_Q^*(\frac{3}{2}^-)$, $\Xi_Q^*(\frac{3}{2}^-)$, $\Omega_Q^*(\frac{3}{2}^-)$, $\Xi_{QQ}^*(\frac{3}{2}^-)$ and $\Omega_{QQ}^*(\frac{3}{2}^-)$ by subtracting the contributions from the corresponding $\frac{1}{2}^+$ and $\frac{3}{2}^+$ heavy and doubly heavy baryon states with the QCD sum rules in a systematic way, and make reasonable predictions for their masses. The present predictions can be confronted with the experimental data in the future at the BESIII, PANDA and LHCb, especially the LHCb. Once reasonable values of the pole residues λ_- are obtained, we can take them as basic input parameters and study the relevant hadronic processes with the QCD sum rules.

Acknowledgements

This work is supported by National Natural Science Foundation, Grant Numbers 10775051, 11075053, and Program for New Century Excellent Talents in University, Grant Number NCET-07-0282, and Project Supported by Chinese Universities Scientific Fund.

Appendix

The spectral densities of the $\frac{1}{2}^-$ and $\frac{3}{2}^-$ heavy and doubly heavy baryon states $\Sigma_Q(\frac{1}{2}^-)$, $\Xi'_Q(\frac{1}{2}^-)$, $\Omega_Q(\frac{1}{2}^-)$, $\Xi_{QQ}(\frac{1}{2}^-)$, $\Omega_{QQ}(\frac{1}{2}^-)$, $\Sigma_Q^*(\frac{3}{2}^-)$, $\Xi_Q^*(\frac{3}{2}^-)$, $\Omega_Q^*(\frac{3}{2}^-)$, $\Xi_{QQ}^*(\frac{3}{2}^-)$ and $\Omega_{QQ}^*(\frac{3}{2}^-)$ at the level of quark-gluon degrees of freedom,

$$\begin{aligned}
\rho_{\Sigma_Q}^A(p_0) &= \frac{1}{2}\rho_{\Omega_Q}^A(p_0) \big|_{\text{RP}}, \\
\rho_{\Sigma_Q}^B(p_0) &= \frac{1}{2}\rho_{\Omega_Q}^B(p_0) \big|_{\text{RP}}, \\
\rho_{\Sigma_Q^*}^A(p_0) &= \frac{1}{2}\rho_{\Omega_Q^*}^A(p_0) \big|_{\text{RP}}, \\
\rho_{\Sigma_Q^*}^B(p_0) &= \frac{1}{2}\rho_{\Omega_Q^*}^B(p_0) \big|_{\text{RP}}, \\
\rho_{\Xi_{QQ}}^A(p_0) &= \rho_{\Omega_{QQ}}^A(p_0) \big|_{\text{RP}}, \\
\rho_{\Xi_{QQ}}^B(p_0) &= \rho_{\Omega_{QQ}}^B(p_0) \big|_{\text{RP}}, \\
\rho_{\Xi_{QQ}^*}^A(p_0) &= \rho_{\Omega_{QQ}^*}^A(p_0) \big|_{\text{RP}}, \\
\rho_{\Xi_{QQ}^*}^B(p_0) &= \rho_{\Omega_{QQ}^*}^B(p_0) \big|_{\text{RP}},
\end{aligned} \tag{14}$$

$$\begin{aligned}
\rho_{\Xi'_Q}^A(p_0) &= \frac{p_0}{32\pi^4} \int_{t_i}^1 dt t (1-t)^3 (p_0^2 - \tilde{m}_Q^2) (5p_0^2 - 3\tilde{m}_Q^2) - \frac{p_0 m_s \langle \bar{q}q \rangle}{4\pi^2} \int_{t_i}^1 dt t \\
&+ \frac{m_s \langle \bar{s}s \rangle}{4\pi^2} \int_{t_i}^1 dt t (1-t) \left[3p_0 + \frac{\tilde{m}_Q^2}{2} \delta(p_0 - \tilde{m}_Q) \right] \\
&- \frac{m_s \langle \bar{s}g_s \sigma G s \rangle}{24\pi^2} \int_0^1 dt t \left[1 + \frac{\tilde{m}_Q}{4T} \right] \delta(p_0 - \tilde{m}_Q) \\
&+ \frac{m_s \langle \bar{q}g_s \sigma G q \rangle}{32\pi^2} \delta(p_0 - m_Q) + \frac{\langle \bar{q}q \rangle \langle \bar{s}s \rangle}{6} \delta(p_0 - m_Q) \\
&+ \frac{p_0}{96\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{t_i}^1 dt (4-5t) + \frac{1}{192\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{t_i}^1 dt (1-t) \tilde{m}_Q^2 \delta(p_0 - \tilde{m}_Q) \\
&- \frac{m_Q^2}{288\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_0^1 dt \frac{(1-t)^3}{t^2} \left[1 + \frac{\tilde{m}_Q}{4T} \right] \delta(p_0 - \tilde{m}_Q),
\end{aligned} \tag{15}$$

$$\begin{aligned}
\rho_{\Xi'_Q}^B(p_0) &= \frac{3m_Q}{64\pi^4} \int_{t_i}^1 dt (1-t)^2 (p_0^2 - \tilde{m}_Q^2)^2 - \frac{m_s m_Q \langle \bar{q}q \rangle}{2\pi^2} \int_{t_i}^1 dt + \frac{m_s m_Q \langle \bar{s}s \rangle}{8\pi^2} \int_{t_i}^1 dt \\
&+ \frac{m_s [6\langle \bar{q}g_s \sigma G q \rangle - \langle \bar{s}g_s \sigma G s \rangle]}{96\pi^2} \delta(p_0 - m_Q) + \frac{\langle \bar{q}q \rangle \langle \bar{s}s \rangle}{3} \delta(p_0 - m_Q) \\
&+ \frac{m_Q}{192\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{t_i}^1 dt \left[-3 - 2t + \frac{2}{t^2} \right] \\
&- \frac{m_Q}{384\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_0^1 dt \frac{(1-t)^2}{t} \tilde{m}_Q \delta(p_0 - \tilde{m}_Q),
\end{aligned} \tag{16}$$

$$\begin{aligned}
\rho_{\Omega_Q}^A(p_0) = & \frac{p_0}{16\pi^4} \int_{t_i}^1 dt t (1-t)^3 (p_0^2 - \tilde{m}_Q^2) (5p_0^2 - 3\tilde{m}_Q^2) - \frac{p_0 m_s \langle \bar{s}s \rangle}{\pi^2} \int_{t_i}^1 dt t \\
& + \frac{m_s \langle \bar{s}s \rangle}{\pi^2} \int_{t_i}^1 dt t (1-t) \left[3p_0 + \frac{\tilde{m}_Q^2}{2} \delta(p_0 - \tilde{m}_Q) \right] \\
& - \frac{m_s \langle \bar{s}g_s \sigma G s \rangle}{6\pi^2} \int_0^1 dt t \left[1 + \frac{\tilde{m}_Q}{4T} \right] \delta(p_0 - \tilde{m}_Q) + \frac{\langle \bar{s}s \rangle^2}{3} \delta(p_0 - m_Q) \\
& + \frac{m_s \langle \bar{s}g_s \sigma G s \rangle}{8\pi^2} \delta(p_0 - m_Q) + \frac{p_0}{48\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{t_i}^1 dt (4-5t) \\
& + \frac{1}{96\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{t_i}^1 dt (1-t) \tilde{m}_Q^2 \delta(p_0 - \tilde{m}_Q) \\
& - \frac{m_Q^2}{144\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_0^1 dt \frac{(1-t)^3}{t^2} \left[1 + \frac{p_0}{4T} \right] \delta(p_0 - \tilde{m}_Q), \tag{17}
\end{aligned}$$

$$\begin{aligned}
\rho_{\Omega_Q}^B(p_0) = & \frac{3m_Q}{32\pi^4} \int_{t_i}^1 dt (1-t)^2 (p_0^2 - \tilde{m}_Q^2)^2 - \frac{3m_s m_Q \langle \bar{s}s \rangle}{2\pi^2} \int_{t_i}^1 dt \\
& + \frac{5m_s \langle \bar{s}g_s \sigma G s \rangle}{24\pi^2} \delta(p_0 - m_Q) + \frac{2\langle \bar{s}s \rangle^2}{3} \delta(p_0 - m_Q) \\
& + \frac{m_Q}{96\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{t_i}^1 dt \left[-3 - 2t + \frac{2}{t^2} \right] \\
& - \frac{m_Q}{192\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_0^1 dt \frac{(1-t)^2}{t} \tilde{m}_Q \delta(p_0 - \tilde{m}_Q), \tag{18}
\end{aligned}$$

$$\begin{aligned}
\rho_{\Omega_{QQ}}^A(p_0) = & \frac{3p_0}{8\pi^4} \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta \alpha \beta (1-\alpha-\beta) (p_0^2 - \tilde{m}_Q^2) (5p_0^2 - 3\tilde{m}_Q^2) \\
& + \frac{3m_Q^2 p_0}{8\pi^4} \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta (1-\alpha-\beta) (p_0^2 - \tilde{m}_Q^2) \\
& - \frac{m_Q^2}{24\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta (1-\alpha-\beta) \left[\frac{\alpha}{\beta^2} + \frac{\beta}{\alpha^2} \right] \left[1 + \frac{p_0}{4T} \right] \delta(p_0 - \tilde{m}_Q) \\
& - \frac{m_Q^4}{192\pi^2 p_0 T} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta (1-\alpha-\beta) \left[\frac{1}{\alpha^3} + \frac{1}{\beta^3} \right] \delta(p_0 - \tilde{m}_Q) \\
& + \frac{m_Q^2}{32\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta (1-\alpha-\beta) \left[\frac{1}{\alpha^2} + \frac{1}{\beta^2} \right] \delta(p_0 - \tilde{m}_Q) \\
& + \frac{m_s \langle \bar{s}s \rangle}{4\pi^2} \int_{\alpha_i}^{\alpha_f} d\alpha \alpha (1-\alpha) \left[6p_0 + p_0^2 \delta(p_0 - \tilde{m}_Q) \right] + \frac{m_s m_Q^2 \langle \bar{s}s \rangle}{8\pi^2} \int_{\alpha_i}^{\alpha_f} d\alpha \delta(p_0 - \tilde{m}_Q) \\
& + \frac{1}{32\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta (\alpha + \beta) \left[3p_0 + \frac{\tilde{m}_Q^2}{2} \delta(p_0 - \tilde{m}_Q) \right] \\
& + \frac{m_Q^2}{64\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta \frac{\alpha + \beta}{\alpha\beta} \delta(p_0 - \tilde{m}_Q) \\
& - \frac{m_s \langle \bar{s}g_s \sigma G s \rangle}{4\pi^2} \int_{\alpha_i}^{\alpha_f} d\alpha \alpha (1-\alpha) \left[1 + \frac{3p_0}{8T} + \frac{p_0^2}{16T^2} \right] \delta(p_0 - \tilde{m}_Q), \tag{19}
\end{aligned}$$

$$\begin{aligned}
\rho_{\Omega_{QQ}}^B(p_0) = & \frac{3m_s}{8\pi^4} \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta \alpha (p_0^2 - \tilde{m}_Q^2)(2p_0^2 - \tilde{m}_Q^2) \\
& + \frac{3m_s m_Q^2}{4\pi^4} \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta (p_0^2 - \tilde{m}_Q^2) \\
& - \frac{m_s m_Q^2}{96\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta \left[\frac{\alpha}{\beta^2} + \frac{\beta}{\alpha^2} \right] \left[\frac{1}{\tilde{m}_Q} + \frac{1}{2T} \right] \delta(p_0 - \tilde{m}_Q) \\
& - \frac{m_s m_Q^4}{96\pi^2 p_0^2 T} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta \left[\frac{1}{\alpha^3} + \frac{1}{\beta^3} \right] \delta(p_0 - \tilde{m}_Q) \\
& + \frac{m_s m_Q^2}{16\pi^2 p_0} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta \left[\frac{1}{\alpha^2} + \frac{1}{\beta^2} \right] \delta(p_0 - \tilde{m}_Q) \\
& - \frac{\langle \bar{s}s \rangle}{2\pi^2} \int_{\alpha_i}^{\alpha_f} d\alpha \alpha (1-\alpha) \left[3p_0^2 - 2\tilde{m}_Q^2 \right] - \frac{m_Q^2 \langle \bar{s}s \rangle}{\pi^2} \int_{\alpha_i}^{\alpha_f} d\alpha \\
& - \frac{m_s}{16\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta \left[1 + \frac{\tilde{m}_Q}{4} \delta(p_0 - \tilde{m}_Q) \right] \\
& + \frac{3\langle \bar{s}g_s \sigma Gs \rangle}{4\pi^2} \int_{\alpha_i}^{\alpha_f} d\alpha \alpha (1-\alpha) \left[1 + \left(\frac{p_0}{2} + \frac{p_0^2}{8T} \right) \delta(p_0 - \tilde{m}_Q) \right] \\
& - \frac{\langle \bar{s}g_s \sigma Gs \rangle}{8\pi^2} \int_{\alpha_i}^{\alpha_f} d\alpha \alpha (1-\alpha) \left[1 + \frac{3p_0}{4} \delta(p_0 - \tilde{m}_Q) \right], \tag{20}
\end{aligned}$$

$$\begin{aligned}
\rho_{\Xi_Q^*}^A(p_0) = & \frac{p_0}{128\pi^4} \int_{t_i}^1 dt t (2+t)(1-t)^2 (p_0^2 - \tilde{m}_Q^2)^2 + \frac{p_0 m_s \langle \bar{s}s \rangle}{16\pi^2} \int_{t_i}^1 dt t^2 \\
& - \frac{m_s \langle \bar{q}q \rangle}{8\pi^2} \int_{t_i}^1 dt t + \frac{m_s \langle \bar{s}g_s \sigma Gs \rangle}{192\pi^2} \int_0^1 dt t \delta(p_0 - \tilde{m}_Q) \\
& + \frac{m_s [3\langle \bar{q}g_s \sigma Gq \rangle - \langle \bar{s}g_s \sigma Gs \rangle]}{192\pi^2} \delta(p_0 - m_Q) + \frac{\langle \bar{q}q \rangle \langle \bar{s}s \rangle}{12} \delta(p_0 - m_Q) \\
& - \frac{p_0}{384\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{t_i}^1 dt t (2-t) - \frac{m_Q^2}{2304\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{t_i}^1 dt \frac{t^3 - 3t + 2}{t^2} \delta(p_0 - \tilde{m}_Q) \tag{21} \\
\rho_{\Xi_Q}^B(p_0) = & \frac{m_Q}{128\pi^4} \int_{t_i}^1 dt (2+t)(1-t)^2 (p_0^2 - \tilde{m}_Q^2)^2 + \frac{m_s m_Q \langle \bar{s}s \rangle}{16\pi^2} \int_{t_i}^1 dt t - \frac{m_s m_Q \langle \bar{q}q \rangle}{8\pi^2} \int_{t_i}^1 dt \\
& + \frac{m_s m_Q \langle \bar{s}g_s \sigma Gs \rangle}{192\pi^2 p_0} \int_0^1 dt t \delta(p_0 - \tilde{m}_Q) + \frac{m_s [3\langle \bar{q}g_s \sigma Gq \rangle - \langle \bar{s}g_s \sigma Gs \rangle]}{192\pi^2} \delta(p_0 - m_Q) \\
& + \frac{\langle \bar{q}q \rangle \langle \bar{s}s \rangle}{12} \delta(p_0 - m_Q) - \frac{m_Q}{384\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{t_i}^1 dt (2-t) \\
& + \frac{m_Q}{1152\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{t_i}^1 dt \frac{(1-t)^3 (3t+4)}{t^2} \\
& - \frac{m_Q}{2304\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_0^1 dt \frac{t^3 - 3t + 2}{t} \tilde{m}_Q \delta(p_0 - \tilde{m}_Q), \tag{22}
\end{aligned}$$

$$\begin{aligned}
\rho_{\Omega_Q}^A(p_0) = & \frac{p_0}{64\pi^4} \int_{t_i}^1 dt t(2+t)(1-t)^2(p_0^2 - \tilde{m}_Q^2)^2 - \frac{p_0 m_s \langle \bar{s}s \rangle}{4\pi^2} \int_{t_i}^1 dt t(2-t) \\
& + \frac{m_s \langle \bar{s}g_s \sigma G s \rangle}{48\pi^2} \int_0^1 dt t \delta(p_0 - \tilde{m}_Q) + \frac{m_s \langle \bar{s}g_s \sigma G s \rangle}{24\pi^2} \delta(p_0 - m_Q) + \frac{\langle \bar{s}s \rangle^2}{6} \delta(p_0 - m_Q) \\
& - \frac{p_0}{192\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{t_i}^1 dt t(2-t) + \frac{m_Q^2}{1152\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{t_i}^1 dt \frac{(1-t)^3}{t^2} \delta(p_0 - \tilde{m}_Q) \\
& - \frac{m_Q^2}{384\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_0^1 dt \frac{(1-t)^2}{t^2} \delta(p_0 - \tilde{m}_Q), \tag{23}
\end{aligned}$$

$$\begin{aligned}
\rho_{\Omega_Q^*}^B(p_0) = & \frac{m_Q}{64\pi^4} \int_{t_i}^1 dt(2+t)(1-t)^2(p_0^2 - \tilde{m}_Q^2)^2 - \frac{m_s m_Q \langle \bar{s}s \rangle}{4\pi^2} \int_{t_i}^1 dt(2-t) \\
& + \frac{m_s m_Q \langle \bar{s}g_s \sigma G s \rangle}{48\pi^2 p_0} \int_0^1 dt \delta(p_0 - \tilde{m}_Q) + \frac{m_s \langle \bar{s}g_s \sigma G s \rangle}{24\pi^2} \delta(p_0 - m_Q) + \frac{\langle \bar{s}s \rangle^2}{6} \delta(p_0 - m_Q) \\
& - \frac{m_Q}{192\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{t_i}^1 dt(2-t) + \frac{m_Q}{576\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{t_i}^1 dt \frac{(1-t)^3(3t+4)}{t^2} \\
& - \frac{m_Q}{1152\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_0^1 dt \frac{t^3 - 3t + 2}{t} \tilde{m}_Q \delta(p_0 - \tilde{m}_Q), \tag{24}
\end{aligned}$$

$$\begin{aligned}
\rho_{\Omega_{QQ}^*}^A(p_0) = & \frac{3p_0}{16\pi^4} \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta \alpha \beta (1-\alpha-\beta)(p_0^2 - \tilde{m}_Q^2)(2p_0^2 - \tilde{m}_Q^2) \\
& + \frac{3m_Q^2 p_0}{16\pi^4} \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta (1-\alpha-\beta)(p_0^2 - \tilde{m}_Q^2) \\
& - \frac{m_Q^2}{192\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta (1-\alpha-\beta) \left[\frac{\alpha}{\beta^2} + \frac{\beta}{\alpha^2} \right] \left[1 + \frac{p_0}{2T} \right] \delta(p_0 - \tilde{m}_Q) \\
& - \frac{m_Q^4}{384\pi^2 p_0 T} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta (1-\alpha-\beta) \left[\frac{1}{\alpha^3} + \frac{1}{\beta^3} \right] \delta(p_0 - \tilde{m}_Q) \\
& + \frac{m_Q^2}{64\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta (1-\alpha-\beta) \left[\frac{1}{\alpha^2} + \frac{1}{\beta^2} \right] \delta(p_0 - \tilde{m}_Q) \\
& + \frac{m_s \langle \bar{s}s \rangle}{4\pi^2} \int_{\alpha_i}^{\alpha_f} d\alpha \alpha (1-\alpha) \left[p_0 + \frac{p_0^2}{4} \delta(p_0 - \tilde{m}_Q) \right] + \frac{m_s m_Q^2 \langle \bar{s}s \rangle}{16\pi^2} \int_{\alpha_i}^{\alpha_f} d\alpha \delta(p_0 - \tilde{m}_Q) \\
& - \frac{1}{48\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta (1-\alpha-\beta) \left[p_0 + \frac{p_0^2}{8} \delta(p_0 - \tilde{m}_Q) \right] \\
& - \frac{m_s \langle \bar{s}g_s \sigma G s \rangle}{32\pi^2} \int_{\alpha_i}^{\alpha_f} d\alpha \alpha (1-\alpha) \left[1 + \frac{2p_0}{3T} + \frac{p_0^2}{6T^2} \right] \delta(p_0 - \tilde{m}_Q), \tag{25}
\end{aligned}$$

$$\begin{aligned}
\rho_{\Omega_{QQ}^*}^B(p_0) = & \frac{3m_s}{32\pi^4} \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta \alpha \beta (p_0^2 - \tilde{m}_Q^2) (3p_0^2 - 2\tilde{m}_Q^2) \\
& + \frac{3m_s m_Q^2}{16\pi^4} \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta (p_0^2 - \tilde{m}_Q^2) \\
& - \frac{m_s m_Q^2}{384\pi^2 T} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta \left[\frac{\alpha}{\beta^2} + \frac{\beta}{\alpha^2} \right] \delta(p_0 - \tilde{m}_Q) \\
& - \frac{m_s m_Q^4}{384\pi^2 p_0^2 T} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta \left[\frac{1}{\alpha^3} + \frac{1}{\beta^3} \right] \delta(p_0 - \tilde{m}_Q) \\
& + \frac{m_s m_Q^2}{64\pi^2 p_0} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta \left[\frac{1}{\alpha^2} + \frac{1}{\beta^2} \right] \delta(p_0 - \tilde{m}_Q) \\
& - \frac{\langle \bar{s}s \rangle}{4\pi^2} \int_{\alpha_i}^{\alpha_f} d\alpha \alpha (1-\alpha) \left[2p_0^2 - \tilde{m}_Q^2 \right] - \frac{m_Q^2 \langle \bar{s}s \rangle}{4\pi^2} \int_{\alpha_i}^{\alpha_f} d\alpha \\
& - \frac{m_s}{64\pi^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_{\alpha_i}^{\alpha_f} d\alpha \int_{\beta_i}^{1-\alpha} d\beta \left[1 + \frac{p_0}{6} \delta(p_0 - \tilde{m}_Q) \right] \\
& + \frac{\langle \bar{s}g_s \sigma G s \rangle}{16\pi^2} \int_{\alpha_i}^{\alpha_f} d\alpha \alpha (1-\alpha) \left[3 + \left(2p_0 + \frac{p_0^2}{2T} \right) \delta(p_0 - \tilde{m}_Q) \right], \quad (26)
\end{aligned}$$

where $\alpha_f = \frac{1+\sqrt{1-4m_Q^2/p_0^2}}{2}$, $\alpha_i = \frac{1-\sqrt{1-4m_Q^2/p_0^2}}{2}$, $\beta_i = \frac{\alpha m_Q^2}{\alpha p_0^2 - m_Q^2}$, $\tilde{m}_Q^2 = \frac{(\alpha+\beta)m_Q^2}{\alpha\beta}$, $\tilde{\tilde{m}}_Q^2 = \frac{m_Q^2}{\alpha(1-\alpha)}$ in the channels $\Xi_{QQ}(\frac{1}{2}^-)$, $\Omega_{QQ}(\frac{1}{2}^-)$, $\Xi_{QQ}^*(\frac{3}{2}^-)$ and $\Omega_{QQ}^*(\frac{3}{2}^-)$; and $\tilde{m}_Q^2 = \frac{m_Q^2}{t}$, $t_i = \frac{m_Q^2}{p_0^2}$ in the channels $\Sigma_Q(\frac{1}{2}^-)$, $\Xi_Q'(\frac{1}{2}^-)$, $\Omega_Q(\frac{1}{2}^-)$, $\Sigma_Q^*(\frac{3}{2}^-)$, $\Xi_Q^*(\frac{3}{2}^-)$ and $\Omega_Q^*(\frac{3}{2}^-)$; the RP denotes the replacements $\langle \bar{s}s \rangle \rightarrow \langle \bar{q}q \rangle$, $\langle \bar{s}g_s \sigma G s \rangle \rightarrow \langle \bar{q}g_s \sigma G q \rangle$ and $m_s \rightarrow 0$.

References

- [1] B. Aubert et al, Phys. Rev. Lett. **97** (2006) 232001.
- [2] K. Nakamura et al, J. Phys. **G37** (2010) 075021.
- [3] V. Abazov et al, Phys. Rev. Lett. **101** (2008) 232002.
- [4] T. Aaltonen et al, Phys. Rev. **D80** (2009) 072003.
- [5] M. Mattson et al, Phys. Rev. Lett. **89**, 112001 (2002).
- [6] A. Ocherashvili et al, Phys. Lett. **B628**, 18 (2005).
- [7] B. Aubert et al, Phys. Rev. **D74**, 011103 (2006).
- [8] R. Chistov et al, Phys. Rev. Lett. **97**, 162001 (2006).
- [9] T. Lesiak, hep-ex/0612042.
- [10] J. L. Rosner, J. Phys. **G34** (2007) S127.
- [11] M. Paulini, arXiv:0906.0808.
- [12] E. Klempt and J. M. Richard, Rev. Mod. Phys. **82** (2010) 1095.

- [13] G. Kane and A. Pierce, "Perspectives On LHC Physics", World Scientific Publishing Company, 2008.
- [14] W. Roberts and M. Pervin, Int. J. Mod. Phys. **A23** (2008) 2817.
- [15] B. Silvestre-Brac, Few Body Syst. **20** (1996) 1.
- [16] S. Capstick and N. Isgur, Phys. Rev. **D34** (1986) 2809.
- [17] S. Migura, D. Merten, B. Metsch and H. Petry, Eur. Phys. J. **A28** (2006) 41.
- [18] D. Ebert, R. N. Faustov and V. O. Galkin, Phys. Lett. **B659** (2008) 612.
- [19] D. Ebert, R. N. Faustov, V. O. Galkin and A. P. Martynenko, Phys. Rev. **D66** (2002) 014008.
- [20] V. V. Kiselev and A. K. Likhoded, Phys. Usp. **45** (2002) 455.
- [21] E. Bagan, M. Chabab and S. Narison, Phys. Lett. **B306** (1993) 350.
- [22] J. R. Zhang and M. Q. Huang, Phys. Rev. **D78** (2008) 094007.
- [23] J. R. Zhang and M. Q. Huang, Phys. Rev. **D78** (2008) 094015.
- [24] H. Garcilazo, J. Vijande and A. Valcarce, J. Phys. **G34** (2007) 961.
- [25] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. **B147** (1979) 385, 448.
- [26] L. J. Reinders, H. Rubinstein and S. Yazaki, Phys. Rept. **127** (1985) 1.
- [27] S. Narison, Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **17** (2002) 1.
- [28] Z. G. Wang, Phys. Lett. **B685** (2010) 59.
- [29] D. Jido, N. Kodama and M. Oka, Phys. Rev. **D54** (1996) 4532.
- [30] Y. Chung, H. G. Dosch, M. Kremer and D. Schall, Nucl. Phys. **B197** (1982) 55.
- [31] E. Bagan, M. Chabab, H. G. Dosch and S. Narison, Phys. Lett. **B301**, 243 (1993).
- [32] Z. G. Wang, Eur. Phys. J. **C68** (2010) 479.
- [33] Z. G. Wang, Eur. Phys. J. **A45** (2010) 267.
- [34] Z. G. Wang, Eur. Phys. J. **C68** (2010) 459.
- [35] J. G. Koerner, D. Pirjol and M. Kraemer, Prog. Part. Nucl. Phys. **33** (1994) 787.
- [36] B. L. Ioffe, Nucl. Phys. **B188** (1981) 317.
- [37] D. Espriu, P. Pascual and R. Tarrach, Nucl. Phys. **B214** (1983) 285.
- [38] B. L. Ioffe, Prog. Part. Nucl. Phys. **56** (2006) 232.
- [39] P. Colangelo and A. Khodjamirian, hep-ph/0010175.
- [40] A. Khodjamirian and R. Ruckl, Adv. Ser. Direct. High Energy Phys. **15** (1998) 345.
- [41] B. Chen, D. X. Wang and A. Zhang, Chin. Phys. **C33** (2009) 1327.
- [42] H. Y. Cheng and C. K. Chua, Phys. Rev. **D75** (2007) 014006.
- [43] D. M. Asner et al, Int. J. Mod. Phys. **A24** (2009) Supp 1.
- [44] M. F. M. Lutz et al, arXiv:0903.3905.